



COLLEGE OF
INTEGRATIVE MEDICINE™

PHONE: 877-841-7241; FAX: 443-327-4763
Info@CollegeofIntegrativeMedicine.org
www.CollegeofIntegrativeMedicine.org
Mail: CIM/Integrative Medicine Health Services, LLC
P.O. Box 407 - Hampstead, MD 21074



Herbal Medicine Approach to Immune Dysfunction

(Part III in a series on Herbal Medicine) (Adapted from The College of Integrative Medicine Module 30 – Clinical Botanical Medicine)

Dr. Wayne Sodano DC, DABCI, DACBN, CFMP, CICP, BCTN

Board Certified Traditional Naturopathy




Image by: Pixabay.com

"Plants have been a central part of traditional medicines to cure topical and systemic infections caused by microbes, in particular bacteria. These preparations form the basis of many wound healing materials in the developing world where the plant is prepared as a crude drug or an extract that is applied topically to improve the healing wound. These preparations may have antimicrobial properties and remove the microbes by an antiseptic mechanism and/or they may promote the ability of the wound to repair itself by stimulating cellular growth."ⁱⁱ
"There are many reasons why plants are available source of antimicrobial natural products and the most fundamental reason is that they contain intrinsically antimicrobial compounds such as carvacrol from thyme (Thymus vulgaris) which is a monoterpene and is present in the essential oil of this species."ⁱⁱⁱ

Herbal Strategies (i.e. phytotherapeutics) to treat infectious diseases includeⁱⁱⁱ:

1. Minor to moderate acute infections of the respiratory, urinary and gastrointestinal mucosa
2. Minor to moderate systemic infections especially when accompanied by lymphadenopathy
3. Topical bacterial and fungal infections
4. Minor to moderate febrile infections
5. Minor to moderate chronic viral, bacterial and fungal infections
6. Management of refractory cases of chronic viral, bacterial and fungal infections especially accompanied by lowered immune resistance.

Herbal Medicine Approach to Immune Dysfunction

(Part III in a series on Herbal Medicine) - (Adapted from The College of Integrative Medicine Module 30 – Clinical Botanical Medicine)

Dr. Wayne Sodano DC, DABCI, DACBN, CFMP, BCTN

Antibacterial / Antibiotic Agents



Indications

Systemic or topical infection.



Contraindications

Serious infections, such as bacterial meningitis; specific to herb.



Application

Best taken before or with meals; medium to long-term application is recommended for extended benefits; support intestinal flora with probiotics.

Primary Herbal Medicine(s)

- o Allium sativum (Garlic)
- o Berberis vulgaris (Barberry)
- o Echinacea spp (Echinacea)
- o Hydrastis Canadensis (Goldenseal)
- o Salvia officinalis (Sage)
- o Scutellaria baicalensis (Baical skullcap)
- o Thymus vulgaris (Thyme)
- o Commiphora molmol (myrrh)



Salvia officinalis



Echinacea spp




Thymus vulgaris

Adult Dosage Consideration

- o Barberry > 15 to 40 ml/week (1:2 liquid); echinacea > 25 to 60 ml/week (1:2 liquid) or 2.5 to 5.0 g/day (tablet).
- o Garlic > 40 to 80 ml/week (1:1 fresh plant liquid) Allicin-releasing enteric-coated tablets containing 6 to 18 mg of allicin per day.
- o Golden seal > 15 to 30 ml/week (1:3 liquid) or 1.0 to 2.0 g/day (tablet).
- o Myrrh > 10 to 30 ml/week (1:5 liquid).
- o Sage > 15 to 30 ml/week 91:2 liquid) or 0.9 to 1.8 g/day (tablet).
- o Thyme > 15 to 40 ml/week (1:2 liquid) or 40 to 60 mg/day (tablet).
- o Baical skullcap > 30 to 60 ml/week (1:2 liquid) or 2.4 to 3.2 g/day (Tablet).

Antifungal Agents



Indications

Fungal infections (systemic or topical).



Contraindications

Specific to herb.



Application

Best taken before meals; medium to long term.

Herbal Medicine Approach to Immune Dysfunction

(Part III in a series on Herbal Medicine) - (Adapted from The College of Integrative Medicine Module 30 – Clinical Botanical Medicine)

Dr. Wayne Sodano DC, DABCI, DACBN, CFMP, BCTN

Primary Herbal Medicine(s)

- o Allium sativum (Garlic)
- o Azadirachta indica (Neem leaf)
- o Tabebuia avellanedae (Pau d'arco)
- o Thuja occidentalis (Thuja) Thyme vulgaris (thyme).
- o Topical - Calendula officinalis (Calendula)
- o Melaleuca alternifolia (Tea tree oil)

Adult Dosage Consideration

- o Garlic > 40 to 80 ml/week (1:1 fresh plant liquid) Allicin-releasing enteric-coated tablets containing 6 to 18 mg of allicin per day; pau d'arco > 25 to 50 ml/week (1:2 liquid) or 1.5 to 2.5 g/day (tablet)
- o Thyme > 15 to 40 ml/week (1:2 liquid) or 40 to 60 mg/day (tablet)
- o Thuja > 15 to 30 ml/week (1:5 liquid) or 400 to 800 mg/day (tablet)

Antiviral Agents



Indications

Suspected mild to moderate viral infections.



Contraindications

Specific to herb.



Application

Best taken before meals; medium to long term.

Primary Herbal Medicine(s)

- o Azadirachta indica (Neem leaf)
- o Hypericum perforatum (St John's wort)
- o Melissa officinalis (Lemon balm)
- o Thuja occidentalis (Thuja)

Adult Dosage Consideration

- o Thuja > 15 to 30 ml/week (1:5 liquid) or 400 to 800 mg/day (tablet)
- o Lemon balm > 20 to 40 ml/week (1:2 liquid)
- o St John's wort > 15 to 40 ml/week (1:2 liquid) or 3.6 to 7.2 g/day (tablet)

Immune Modulators

Modulate or balance immune system activity



Indications

recurrent infectious disease due to weakened immune system; weakened immune system



Contraindications

Specific to herb; individuals receiving immunosuppressant therapy



Application

Best taken before meals; medium to long term.

Herbal Medicine Approach to Immune Dysfunction

(Part III in a series on Herbal Medicine) - (Adapted from The College of Integrative Medicine Module 30 – Clinical Botanical Medicine)

Dr. Wayne Sodano DC, DABCI, DACBN, CFMP, BCTN

Primary Herbal Medicine(s)	Adult Dosage Consideration
<ul style="list-style-type: none">o Echinacea spp. (Echinacea root)o Eleutherococcus senticosus (Siberian ginseng)o Panax ginseng (Korean ginseng)o Withania somnifera (withania, Ashwagandha)	<ul style="list-style-type: none">o Echinacea root > 25 to 60 ml/week (1:2 liquid) or 2.5 to 5.0 g/day (tablet).o Korean ginseng > 7 to 40 ml/week (1:2 liquid) or 300 to 500 mg/day (tblmg/day (tablet).o Siberian ginseng > 15 to 55 ml/week (1:2 liquid) or 2.5 to 6.3 g/day (tablet).o Withania (Ashwagandha) > 20 to 50 ml/week (1:1 liquid) or 2.9 to 4.8 g/day (tablet)

Immune Suppressants

Dampen immune response (for autoimmune diseases, hay fever, asthma, eczema, urticaria, chronic skin disorders)



Indications

Over active immune system



Contraindications

specific to herb; caution with weakened immune systems



Application

Best taken before meals; medium to long term.

Primary Herbal Medicine(s)	Adult Dosage Consideration
<ul style="list-style-type: none">o Albizia lebbek (Albizia)o Hemidesmus indicus (Hemidesmus),o Tylophora indica (Tylophora)	<ul style="list-style-type: none">o Albizia > 25 to 60 ml/week (1:2 liquid) or 2.4 to 4.0 g/day (tablet)o Hemidesmus > 25 to 60 ml/week 91;2 liquid) or 1.5 to 2.5 g/day (tablet)o Tylophora > 5 to 15 ml/week (1:5 liquid) for the first 10 to 14 days of each month - dispense separately from other liquid herbs



Albizia



Hemidesmus



Tylophora

Herbal Medicine Approach to Immune Dysfunction

(Part III in a series on Herbal Medicine) - (Adapted from The College of Integrative Medicine Module 30 – Clinical Botanical Medicine)

Dr. Wayne Sodano DC, DABCI, DACBN, CFMP, BCTN

Lymphatics

Herbal medicines that improve lymphatic flow and increase lymphatic detoxification



Indications

Lymphatic congestion / stagnation; detoxification



Contraindications

Specific to herb; caution with individuals with lymphatic cancer



Application

Best taken before meals; medium to long term.

Primary Herbal Medicine(s)

- o Baptista tinctoria (Wild indigo)
- o Calendula officinalis (Calendula)
- o Echinacea ssp. (Echinacea)
- o Iris versicolor (Blue flag)
- o Galium aparine (Cleavers)
- o Phytolacca decandra (Poke root)

Adult Dosage Consideration

- o Baptista > 15 to 40 mL/week (1:2 liquid)
- o Calendula > 10 to 30 ml/week (1:2 liquid); blue flag > 20 to 40 ml/week (1:2 liquid)
- o Clivers (Galium aparine) > 25 to 50 ml/week (1:2 liquid) or 1.0 to 2.0 g/day (tablet)
- o Echinacea > 25 to 60 ml/week (1:2 liquid) or 2.5 to 5.0 g/day (tablet)
- o Poke root > 1 to 5 ml/week (1:5 tincture)



Baptista tinctoria



Candula officinalis



Iris versicolor



Galium aparine



Echinacea spp



Phytolacca decandra

Poor immunity and recurrent infections^{iv}

Individuals with poor immunity and/or recurrent infections should receive treatment selected from the following groups:

1. Immune-enhancing herbs:

- o Echinacea
- o Astragalus *
- o Picrorrhiza *
- o Andrographis
- o Phytolacca

*Astragalus should not be prescribed during acute episodes and Picrorrhiza and Astragalus should not be prescribed if the patient is constitutionally cold.

Herbal Medicine Approach to Immune Dysfunction

(Part III in a series on Herbal Medicine) - (Adapted from The College of Integrative Medicine Module 30 – Clinical Botanical Medicine)

Dr. Wayne Sodano DC, DABCI, DACBN, CFMP, BCTN

2. Tonic and adaptogenic herbs:

- o Panax*
- o Eleutherococcus*
- o Withania

Panax and Eleutherococcus should not be

3. Bitter herbs:

- o Artemisia absinthium - Especially where the patient appears anemic or undernourished. Exercise caution if the patient is also constitutionally cold or counter the cooling effect with warming herbs.

Sample Echinacea Protocol

Echinacea alone, either the root of *E. angustifolia* or *E. purpurea*, has helped countless patients with poor immunity in doses equivalent to 2.5 - 7.5 g/day (5 - 15 ml of a 1:2 preparation)



Root of *E. Angustifolia*



E. Purpurea



Iris versicolor



1. Take 5 ml dose each day (2.5 g) as a maintenance dose (take this twice this dose for maintenance if immunity is poor.)
2. If infection threatens, double or triple the daily maintenance dose until the treat passes.
3. If infection takes hold, maintain the higher dose until the infection is completely gone and then return to normal daily dose.

References

- ⁱ Heinrich M, Barnes J, Gibbons S, Williamson EM. *Fundamentals of Pharmacognosy and Phytotherapy*. 2nd Ed. Edinburgh: Churchill Livingstone; 2012. p. 252.
- ⁱⁱ *Ibid.*
- ⁱⁱⁱ Mills S, Bone K. *Principles and Practice of Phytotherapy; Modern Herbal Medicine*. Edinburgh: Churchill Livingstone; 2000. p. 140.
- ^{iv} *Ibid.* p. 142.



LIMITS OF LIABILITY & DISCLAIMER OF WARRANTY. The material provided herein is provided for the purpose of education through research of reputable sources within the healthcare industry. The author(s) do not hold liability for any misconception or misuse of the information provided. The material is not meant to be a comprehensive source for the topic covered, and is not intended as a substitute for medical diagnosis, treatment, or medical counseling. Information contained herein should not be construed as a claim or representation that any treatment, process or interpretation mentioned constitutes a cure, palliative, or ameliorative, nor is it intended to supplement the practitioner's knowledge of their patient, and should be considered as adjunctive and support to other diagnostic medical procedures. This material contains elements protected under International and Federal Copyright laws and treaties. Any unauthorized reprint or use of this material is prohibited.